References:

- Ashok Kumar B. S., Lakshman K., Jayaveea K. N., Shekhar D. S., Khan S., Thippeswamy B. S., Veerapur V. P., (2002). Antidiabetic, antihyperlipidemic and antioxidant activities of methanolic extract of Amaranthus viridis Linn in alloxan induced diabetic rats. Experimental and Toxicologic Pathology, 64: 75– 79.

- Bairy K. L., Adiga S., Meharban A., Punita I. S. R., (2010). Hypoglycemic effect of aqueous extract of trichosanthes dioica in normal and diabetic rat. Int J Dia Dev Ctries., 30(1): 38-42.

- Barik R., Jain S., Owatra D., joshi M., Tripathi G. S., Goyal R., (2008). Antidiabetic activity of aqueous root extract of Inchnocarpus *frutescens* in streptozotocin-nicotinamide indiced tipe-2 diabetes. Indian J pharmacol, 40 (1):19-22.

- Bhowmik A., Khan L. A., Akhtar M., Rokheya B., (2009). Studies on the antidiabetic effects of Mangifera indica stem-barks and leaves on nondiabetic, type 1 and type 2 diabetic model rats. Bangladesh J Pharmacol, 4: 110-114.

- Chan, J. C. N., Yeung, V. T. F., Cockram, C. S., (1996). Diabetes Mellitus - Epidemiology and Pathogenesis. Hong Kong Practitioner; 18: 270-279.

- Dewanjee, S., Bose, S. K., Sahu, R., Mandal, S. C., (2008). Antidiabetic effect of matured fruits of *Diopsyros peregrine* in alloxan induced diabetic rats. International Journal of green pharmacy, p. 95-99.

- Gandhipuram, P. S., Palanisamy, A., kumar, D. S., and Subramanian, S. P., (2006). Antidiabetic activity of fruits of Terminalia chebula on streptozotocin induced diabetic rats. Journal of health science, 52(3):283-291.

- Gonzalez-Trujano M.E., Carrera D., Venturs-martinez R., Cedillo-Portugal E., Navarrete A., (2006). Neuropharmacological profile of an ethanol extract of *Ruta chalepensis* L. in mice. Journal of Ethnopharmacology 106: 129–135.

- Hussain A., Hydrie M. Z. I., Claussen B., Asghar S., (2010). Type 2 Diabetes and obesity: A review. Journal of Diabetology, 2(1): 1-7.

- Ignacimuth S., Daisy P., Jasmine R., Murugan E., (2009). A novelSteroid1 from Elephantopus scaber L. an Ethnomedicinalplant with antidiabeticactivity. Phytomedicine 16: 252–257.
- Jangra M., Sharma S., Kumar M., (2010). Evaluation of antihyperglycemic activity of *dodonaea viscosa* leaves in normal and stzdiabetic rats. Int J Pharm Pharm Sci, 3(3): 69-74.
- Jarald E. E., Bangar O. P., Asghar J., Ahmed S., (2009). Antidiabetic activity of polyherbal formulation. International journal of green pharmacy, july-sept: 211-214.
- Ji S. K., Jung B. J., Chang Won Choi and Sei Chang Kim, (2006). Hypoglycemic and Antihyperlipidemic Effect of Four Korean Medicinal Plants in Alloxan Induced Diabetic Rats. American Journal of Biochemistry and Biotechnology, 2 (4): 154-160.
- Kannur D. M., Hukkeri V. M., Akki A. S., (2006). Antidiabetic activity of Caesalpinia bonducella seed extracts in rats. Fitoterapia, 77: 546–549.
- Karthikeyan A., Siva G., Rex D. A. P., Chidambaranathan N., (2008). antihyperglycemic activity of ethanolic seed extract of *vernonia anthelminticum* Willd. International journal of green pharmacy, oct-dec: 209-211.
- Kim, J. D., Kang, S. M., Park, M.Y., Jung, T. Y., Choi, H. Y. and Ku, S. K., (2007). antidiabetic activity of Dangnyosoka, a Chinese herbal medicine, in diabetic rats. Biosci. Biotech. Biochem, 71(6):1527-1534.
- Kirtikar, K. R. and Basu, B. D., (1991). Indian medicinal plants, Vol-II, 5th ed. Delhi: periodical experts book agency, 946-947.
- Kumar A., Llavarasan R., deekaraman R., Aravindan P., Padmanabhan N., Krishnan M. R. V., (2008). Anti-diabetic activity of *Syzygium cumini* and its isolated compound against streptozotocin-induced diabetic rats. J. Med. Plant. Res., 2 (9): 246-249.
- Kurihara H., Bao L., Yao X., Xu J., Guo X., Liu H., (2009). Effects of Pithecellobium clypearia Benth extract and its main components on inflammation and allergy. Fitoterapia, 80: 349–353.

- Lenzen S., (2008). The mechanisms of alloxan- and streptozotocin-induced diabetes. Diabetologia, 5:216–226.
- Mankil J., Park M., Lee h. C., Kang Y., Kim S. K., (2006). Antidiabetic Agents from Medicinal Plants. Current Medicinal Chemistry, 13: 1203-1218.
- Md. Salahuddin and Jalalpure S. S., (2010). Antidiabetic activity of aqueous fruit extracts of Cucumis trigonus Roxb in streptozotocin-induced-diabetic rats. Journal of Ethnopharmacology, 127: 565–567.
- Mohire N. C., Salunkhe V. R., Bhise S. B., Yadav A. V., (2007). Cardio tonic activity of the extract of heartwood of *Pterocarpus marsupium.* Indian J of experimental biology, 45: 532-537.
- Mustafa A. and David E. L.; (2011). Diabetes, oxidative stress and physical exercise. Journal of Sports Science and Medicine 1: 1-14.
- Nagappa A. N., Thakurdesai P. N., Rao V. N., Singh J., (2003). Antidiabetic activity of *Terminalia catappa Linn* fruits. Journal of Ethnopharmacology 88: 45–50.
- Naik P. R. and Mohammadi J. et al; (2008). Evaluation of hypoglycemic effect of Morus alba in an animal model. Indian J Pharmacol, 40 (1): 15-18.
- Orhan N., Asnan M., Orhan D. D., Ergun F., Yesilada E., (2006). In-vivo assessment of antidiabetic and antioxidant activities of grapevine leaves (*Vitis vinifera*) in diabetic rats. Journal of Ethnopharmacology, 108: 280–286.
- Ozturk Y., Altan V. M., Ari N., (1998). Diabetic Complications in Experimental Models. Tr. J. of Medical Sciences, 22: 331-341.
- Pathak A. K. and Argal A., (2006). CNS activity of *Calotropis gigantea* roots. Journal of Ethnopharmacology 106: 142–145.
- Rajendran R., Suseela L., Meemakshi S. R., Saleem B. N., (2008). Cardiac stimulant activity of bark and wood of Premna serratifolia. Bangladesh J Pharmacol; 3: 107-113.
- Rao C. A., Rao B. K., sudarshan P. R., Rajashekhar M. D., (2003). Antidiabetic activity of *Terminalia pallida* fruit in alloxan induced diabetic rats. Journal of Ethnopharmacology, 85: 169–172.

- Rao M. U., Sreenivasulu M., Chengaiah., Jaganmohan reddy k., (2010). Herbal Medicines for Diabetes Mellitus: A review. Int.J. PharmTech Res; 2(3): 1883-1892.
- Rauter A. P., Martins A., Ferreira J., Borges C., Justino J., Silva S. V., Goulart M., Rodrigues J. A., Pinto J. A., (2009). Bioactivity studies and chemical profile of the antidiabetic plant *Genista tenera*, Journal of Ethnopharmacology, 122: 384–393.
- Saha B. K., Bhuiyan n. H., Mazumder K., Farmuzul Haque K. M., (2009). Hypoglycemic activity of Lagerstroemia speciosa L. extract on streptozotocin-induced diabetic rat: Underlying mechanism of action, Bangladesh J Pharmacol, 4: 79-83.
- Satyanarayan, U., (2004). Biochemistry, 2nd ed. Kolkata: book and allied Pvt. Ltd. P. 251-290.
- Shanmugakumar, S. D., Amerjothy, S., Balakrishna, K. and Soo-un Kim, (2008). Isolation, Characterization and in-vitro evaluation of Kaempferol-3-o-alfa-l-Rhamanopyranoside from Pithecellobium dulce benth. against Mycobacterium tuberculosis and its associated secondary infections, Pharmacognosy Magazine, 4(15):59-S64.
- Shanmugakumar, S.D., Amerjothy, S. and Balakrishna, K, (2006). Research Article Pharmacognostical, Antibacterial and Antifungal potentials of the leaf extracts of *Pithecellobium dulce* Benth. Pharmacognosy Magazine, 2(7):163-166.
- Sundarrajan T., Raj Kumar T., Udhaykumar E., Arunachalam G., (2010). Hypolipidemic activity of *Pithecellobium Dulce* Bench. In Triton Wr-1339 Induced Hyperlipidemic rats. International Journal of Chemical and Pharmaceutical Sciences, 2(1): 50-53.

- Szkudelski T.; (2001). The Mechanism of Alloxan and Streptozotocin Action in B Cells of the Rat Pancreas. Physiol. Res, 50: 536-546.
- Watal G., Singh S. K, Kesari A. N., Gupta R. K., Jaiswal D., (2007). Assessment of antidiabetic potential of *Cynodon dactylon* extract in streptozotocin diabetic rats. Journal of Ethnopharmacology 114: 174–179.
- World Health Organization, (1999) Definition, Diagnosisand Classification of Diabetes Mellitus and its Complications- report of a WHO consultation. Geneva. - World Health Organization.